



IRF8-dependent DCs play a key role in the regulation of CD8 T cell responses to epithelial-derived antigen in the steady state but not in inflammation

Joeris, Thorsten

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Joeris, T. (2016). *IRF8-dependent DCs play a key role in the regulation of CD8 T cell responses to epithelial-derived antigen in the steady state but not in inflammation*. Abstract from 10th European Mucosal Immunology Group meeting, Copenhagen, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Along the process of epithelial self-renewal, antigens derived from apoptotic intestinal epithelial cells (IECs) are taken up by antigen presenting cells (APCs), transported to gut-draining lymph nodes and cross-presented to CD8 T cells. In steady state, rapid tolerization of CD8 T cells reactive towards epithelial-derived antigens is crucial to maintain tissue homeostasis. Since IRF8-dependent dendritic cells (IRF8-DCs) have superior cross-presenting capabilities, we aimed to investigate their role in this process. IFABP-tOva mice, expressing the model-antigen Ovalbumin (Ova) in IECs, were used as recipients to set up chimeras using either CD11c-cre.Irf8^{fl/fl} bone marrow, which cannot generate IRF8-DCs, or cre-negative Irf8^{fl/fl} control bone marrow. Whereas transfer of Ova-specific CD8 T cells (OT-I cells) to control chimeras resulted in their rapid tolerization, OT-I cells transferred to CD11c-cre.Irf8^{fl/fl} chimeras spontaneously developed into cytotoxic effector T cells (CTL), causing epithelial destruction and intestinal inflammation. However, when the adjuvant R848 was applied in addition to OT-I transfer, inflammation was triggered in both, CD11c-cre.Irf8^{fl/fl} and control chimeras. This demonstrates that IRF8-DCs are crucial for the rapid tolerization of CD8 T cells reactive towards epithelial-derived antigen in steady state, but are not essential for the induction of CTLs in an inflammatory setting such as found in infection.